arXiv:0707.3387v1 [gr-qc] 23 Jul 2007

Note on Solving for the Dynamics of the Universe

R.F.L. Holanda*
IAG, Universidade de São Paulo, 05508-900 São Paulo, SP, Brasil and
IAG, Universidade de São Paulo, 05508-900 São Paulo, SP, Brasil

Lima (2001)has proposed an alternative procedure to solve the Friedman-Robertson-Walker (FRW) cosmological equation of simple fluid with equation of state $p = (\gamma - 1)\rho$. The basic result of such paper was to reduce the FRW equations to the the form of an oscillator equation. In this note we generalize such method for multidimentional FRW cosmologies.

Recently, Lima [1] rediscussed the cosmological solutions for a relativistic simple fluid in the framework of FRW models. In such paper the cosmological equation of simple fluid with equation of state $p = (\gamma - 1)\rho$ was reduced to a oscillator equation form. The possible dynamic behaviors of the FRW equations were naturally recovered in such reduction. In this note we generalize the method for multidimentional FRW cosmologies.

The spacetime metric for multidimentional FRW cosmologies can take the following form [3] ($c = 1$)

$$ds^2 = dt^2 - a^2(t)\left(1 + \frac{kr^2}{4}\right)^{-1}\delta_{ij}dx^i dx^j \tag{1}$$

where $r^2 = \sum_i x_i^i$ e i, j=1, 2, 3...D.

In such background, the Einstein field equations for a relativistic simple fluid and the conservation of energy can be written as

$$\frac{D(D-1)}{2}\left[\frac{\dot{a}^2}{a^2} + \frac{k}{a^2}\right] = 8\pi G_D \rho + \Lambda \tag{2}$$

$$\frac{(D-1)\ddot{a}}{a} + \frac{(D-1)(D-2)}{2}\left[\frac{\dot{a}^2}{a^2} + \frac{k}{a^2}\right] = -8\pi G_D p + \Lambda \tag{3}$$

$$\dot{\rho} + D(\rho + p)H = 0 \tag{4}$$

where ρ and p are the energy density and pressure, respectively.

In the cosmological framework, it is usually assumed that the matter content obeys the barotropic γ-law equation of state

$$p = (\gamma - 1)\rho\ , \qquad \gamma \in [0, 2] \quad . \tag{5}$$

it is interesting to see that the space dimensionality multiplies directly the parameter γ in the energy density equation as

$$\rho = \rho_0\left(\frac{a_0}{a}\right)^{D\gamma} \tag{6}$$

The cosmic dynamics is determined by combining the above set of equations. In principle, the corresponding dynamic behavior must be heavily dependent on the choice of the three free parameters: (i) The curvature parameter K, (ii) the equation of state parameter γ, and (iii) the spatial dimension D. As one may check, the evolution of the scale function is driven by the second order differential equation

$$a\ddot{a} + \Delta_D \dot{a}^2 + \Delta_D k = 0 \tag{7}$$

and

$$\Delta_D = \left(\frac{D\gamma - 2}{2}\right) \tag{8}$$

It is possible to extend for the case of D-dimensions spatial the method solution proposed by Lima [1]. This can be accomplished by using the conformal time η, instead of the cosmological or physical time [$dt = a(\eta)d\eta$]. In this case the equation of motion (7) takes the form below:

$$aa^{''} + (\Delta_D - 1)a^{'2} + \Delta_D k a^2 = 0 \tag{9}$$

We now employ the auxiliary factor

$$Z(\eta) = lna \ \text{ if } \ \Delta_D = 0 \tag{10}$$

$$Z(\eta) = a^{\Delta_D} \ \text{ if } \ \Delta_D \neq 0 \tag{11}$$

to obtain, respectively,

$$Z^{''} = 0 \ \text{ if } \ \Delta_D = 0 \tag{12}$$

$$Z^{''} + k\Delta_D^2 Z = 0 \ \text{ if } \ \Delta_D \neq 0 \tag{13}$$

Although we are treating FRW cosmologies in D-dimentional space, the equations (12) and (13) are identical to found by Lima for the 3-dimentional case. The eq. (13) stays describing the classical motion of a particle

---

*Electronic address: Holanda@astro.iag.usp.br

subject to a linear force. This force is of restoring or repulsive type depending only on the sign of the curvature parameter. Closed models (k=1) are, for any value of $\Delta_D \neq 0$, analogous to harmonic simple oscillators (HSO).

It should be stressed that the above method based on the transforming equations (10) and (11) is also convenient when new ingredients are considered such as the presence of a cosmological Λ term. The second-order differential equation (7) takes the form

$$a\ddot{a} + \Delta_D \dot{a}^2 + \Delta_D k = \frac{\Lambda \gamma a^2}{D-1} \tag{14}$$

and

$$\Delta_D = \left[\frac{D\gamma - 2}{2}\right]$$

As an advanced exercise, one may show that the generalized equation of motion for Λ models in terms of the auxiliary scale factor is

$$Z^{''} + \Delta^2 k Z = \frac{2\Delta(\Delta+1)\Lambda Z^{\frac{\Delta+2}{\Delta}}}{D(D-1)} \tag{15}$$

When $D \longmapsto 3$ the Lima's results are recovered.

Acknowledgments

This work was partially supported by the Capes (Brazilian research agency). The author is grateful to Prof. J.A.S. Lima for helpful discussions.

[1] J.A.s. Lima,“ Note on Solving for the dynamics of the universe”, Am.J.Phys. 69, 1245-1247,(2001)

[2] V. Faraoni, “Solving for the dynamics of the universe”, Am. J. Phys. 67 (8), 732 (1999).

[3] F.R. Tangherline,“ Dimensionality of space and the pulsating Universe”, N. Cim. 91B, 209,(1986)

[4] M. J. D. Assad and J. A. S. Lima, “General and unified solution for perfect fluid homogeneous and isotropic cosmological models”, Gen. Rel. Grav. 20, 527 (1988).

[5] L. D. Landau and E. M. Lifshitz, The Classical Theory of Fields, (Pergamon, Oxford 1989), pp. 363-367.

[6] R. D’Inverno, Introducing Einstein’s Relativity (Clarendon, Oxford, 1992) p. 322.

[7] Ya. B. Zeldovich and I. D. Novikov, Stars and Relativity (Dover publication, New York, 1996) pp. 186-187.

[8] J. A. S. Lima, Notes in Cosmology 1990 (in Portuguese, unpublished).

[9] J. A. S. Lima, A. M. Moreira and J. Santos, “Particle-like description for FRW cosmologies”, Gen. Rel. Grav. 30, 425 (1998).